

Reductionist synthetic community approaches in root microbiome research

Yong-Xin Liu^{1,2,3}, Yuan Qin^{1,2,3,4} and Yang Bai^{1,2,3,4}



Synthetic community (SynCom) approaches can provide functional and mechanistic insights into how plants regulate their microbiomes, and how the microbiome in turn influences plant growth and health. Microbial cultivation and reconstruction play pivotal roles in this process, which enables researchers to reproducibly investigate the interactions between plants and a major proportion of plant-associated microbes in controlled laboratory conditions. Here, we summarize the emergence, current achievements, and future opportunities for using SynCom experiments in plant microbiome research, with a focus on plant root-associated bacteria.

Addresses

¹ State Key Laboratory of Plant Genomics, Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing 100101, China

² CAS Center for Excellence in Biotic Interactions, University of Chinese Academy of Sciences, Beijing 100049, China

³ CAS-JIC Centre of Excellence for Plant and Microbial Science (CEPAMS), Institute of Genetics and Developmental Biology, Chinese Academy of Sciences (CAS), Beijing 100101, China

⁴ College of Advanced Agricultural Sciences, University of Chinese Academy of Sciences, Beijing 100049, China

Corresponding author: Bai, Yang (ybai@genetics.ac.cn)

Current Opinion in Microbiology 2019, 49:97–102

This review comes from a themed issue on **Environmental microbiology**

Edited by **Roeland Berendsen** and **Klaus Schlaeppli**

<https://doi.org/10.1016/j.mib.2019.10.010>

1369-5274/© 2019 Published by Elsevier Ltd.

Introduction

A distinctive microbiome assembles on plant roots, with microbial members originating from soil environments [1^{••},2,3^{••}]. These microbes interact with plants throughout their lifecycle [4,5], influencing plant growth [6,7^{••}], nutrient uptake [8^{••},9^{••}], and disease resistance [10^{••},11,12,13[•]]. Over the past decade, root microbiome compositions have been extensively investigated in model [1^{••},3^{••},14] and crop plants [10^{••},15^{••},16–19]. In addition, researchers have cultivated bacteria and established stocks from the microbiomes of a variety of plants,

including *Arabidopsis thaliana*, rice (*Oryza sativa*), maize (*Zea mays*), tomato (*Solanum lycopersicum*), potato (*Solanum tuberosum*), clover (*Trifolium pratense*), and sugarcane (*Saccharum* sp.) [8^{••},10^{••},18,20^{••},21^{••},22,23]. The ultimate goal of culturing these bacteria is to examine the functions and mechanisms of the interactions between root microbiomes and host plants. Here, we discuss the key challenges of using experimental synthetic communities (SynComs), namely microbiome cultivation and reconstitution.

From holism to reductionism

Research into the interactions between root microbiomes and host plants connects the fields of microbial ecology and plant molecular biology. Ecological studies typically investigate the root microbiome using the holistic approaches, which is critical to reveal the root microbiome status in natural environments. By contrast, studies using the reductionist approaches try to avoid uncontrollable variables, breaking down the interactions of the root microbiome with the plant host into experimentally controllable factors (cultivated microbes, nutrients, plant genotype, and so on) to understand their functions and mechanisms [24[•]] (Figure 1). Two seminal works pioneered the reductionist concept in root microbiome research [1^{••},3^{••}]. The authors of these studies grew *Arabidopsis* plants in natural soils under controlled conditions to avoid differences in wind, rain, and temperature while allowing plants to assemble a root microbiome from native soil microbial communities (Figure 1a,b). Using a large number of biological and technical replicates, the two studies concluded that *Arabidopsis* plants assemble a distinctive root microbiome in comparison with the microbiome of unplanted soils. In addition, Bulgarelli *et al.* demonstrated that the major assembly patterns of root microbiomes in natural soils under laboratory conditions were similar to the root microbiomes in natural field sites [1^{••}].

Investigating the assembly of the root microbiome from a natural soil inoculum under controlled conditions became popular for root microbiome studies in species such as rice, *Arabidopsis* and its relatives [15^{••},25^{••},26[•]] (Figure 1b). Reducing environmental variation is critical for meaningful comparative descriptions of microbiomes; however, in order to experimentally manipulate microbiomes, it is essential to dissect the interactions between the root microbiome and the host plants into experimentally controllable factors, such as microbes, plant genotypes, nutrients, and growth conditions (Figure 1c). This requires that researchers cultivate specific microbes [27].

Figure 1




Illustration of microbiota research using field conditions, controlled growth chambers, and synthetic community (SynCom) systems. **(a)** Field experiments are used to explore the assembly and variation of the root microbiome in natural conditions. Environmental factors, such as temperature, wind, rain, microbiota, and soil nutrients, cannot be controlled. **(b)** Experiments using natural soils in controlled growth chambers can be used to avoid environmental variations and provide reproducible results. The microbiota members and soil nutrients cannot be dissected using this technique. **(c)** SynCom approaches use cultivated microbes to provide unique opportunities to obtain fully reproducible results and to investigate the function of the root microbiome in plant fitness.

Cultivating and dissecting the root microbiome


Approaches that manipulate the microbiota require extensive collections of microbial cultures and axenic systems [20^{**},28^{**},29^{**}]. Although thousands of cultivated microbes have been isolated and stored in international stock centers [30], these microbes were obtained from various environments or host species, and may have adaptations to local conditions and to interactions with other components of the native microbiome. Therefore, existing isolates may be genotypically and functionally distinct from the microbes that researchers find in their new samples, thus may not be suitable to dissect the interactions between plants and the root microbiome in a given natural soil.

High-throughput bacterial cultivation provides a solution for capturing and manipulating a bacterial community on a large scale (Figure 2a–d). Recently, Bai *et al.* established a pipeline for high-throughput bacterial cultivation and identification, enabling the characterization of 7943 bacterial isolates from roots and leaves of *Arabidopsis* plants grown in natural soils [20^{**}]. This technique involves the use of tryptone yeast extract glucose, Reasoner's 2A, and related media with the limiting dilution method, complemented with cell sorting and colony picking. The bacterial isolates were characterized using a 454-based protocol, which allowed the authors to identify co-cultivated strains. The approach facilitated the cultivation of more than 50% of the bacterial taxa reproducibly present in the *Arabidopsis* root and leaf microbiomes. More recently, Zhang *et al.* used an improved identification method to characterize 13 512 bacterial isolates from rice

roots, covering 70% of the rice root bacterial microbiome members. This new method improved the sequencing depth using Illumina HiSeq2500, and increased accuracy of identification by avoiding chimera formation using two-side barcode labeling and independent library preparation for each isolate [8^{**}]. These methods provide an effective way to dissect a specific root microbiome sample with the advantage of being able to identify co-cultivated bacteria (i.e. bacterial 'isolates' that contain more than one type of bacteria).

To achieve as much diversity as possible, targeted isolation of fastidious bacteria (i.e. bacteria with specific culture requirement not met by the standard media) can complement high-throughput bacterial cultivation. Several approaches based on the 16S rRNA gene sequence or the metagenome can help identify the appropriate specialized isolation media. For example, Oberhardt *et al.* developed a tool to predict the isolation media required using the bacterial 16S rRNA gene sequence [31]. In another example, using a Greengene ID, the BugBase database can be used to obtain detailed bacterial characteristics, such as aerobic or anaerobic growth, which provide guidance for the types of cultivation conditions required [32]. Metagenomic data or bacterial genomes present a second source of information about bacterial carbon utilization and antibiotic resistance genes, which aids the design of isolation media to enriched specific bacteria [33,34]. Guided by metagenomic data, Kwak *et al.* used a marine broth medium containing kanamycin to successfully isolate 22 flavobacteria strains that confer wilt resistance in tomato [10^{**}]. It is important to be careful with the medium preparation

Figure 2



SynCom workflow.

technique, however; for example, Kato *et al.* reported that autoclaving phosphate and agar together generates H_2O_2 , which decreases the resulting colony count [35]. Future cultivation effort should be targeted towards gathering ‘hard-to-get’ taxa (i.e. Acidobacteria, Chloroflexi, Deltaproteobacteria and anaerobic taxa), which ubiquitously exist in root microbiota but not well represented in culture collections.

The first-pass characterization and selection of isolates in SynCom experiments often use the 16S rRNA gene sequence of cultivated strains for cross-referencing the microbiota profiling data (Figure 2e). In microbiota profiling, comparing with operational taxonomic units (OTUs) clustering using 97% 16S rRNA gene sequence similarity, amplicon sequencing variants (ASVs, 100% 16S rDNA sequence similarity) obtained in UNOISE, DADA2, and Deblur pipelines provide a higher resolution. ASVs distinguish single nucleotide polymorphisms in the 16S rRNA gene; this allows a more accurate comparison between the cultivated bacteria and amplicon data [36–38]; however, it should be noted that the 16S rRNA gene does not reveal all genomic and functional variations among bacterial isolates, despite being the most widely used taxonomic marker. For example, Karasov *et al.* demonstrated that *Pseudomonas* isolates belonging to the same OTU (based on 16S rRNA gene sequences) showed clear differentiation at the level of genomic content and disease phenotypes [39]. Selecting bacterial strains with the same 16S rRNA gene sequence but isolated from different roots, representing different

colonization events, will therefore increase the functional diversity of the cultivated root microbiome members [8^{**},20^{**}].

Comparison of axenic systems

SynComs mainly use three axenic systems: agar-based, clay-based, and FlowPot systems, each system with different advantages and disadvantages. An agar system is highly artificial but many factors, such as nutrient levels and biotic and abiotic stresses, are uniformly controlled [40^{**},41,42]. Compared with an agar system, a clay-based system better mimics soil conditions because the texture of clay is similar to that of soil. However, the clay matrix influences the nutrient status, making it difficult to deplete specific nutrients such as nitrogen or phosphate, and the clay lacks organic carbon [8^{**},20^{**},43^{**}]. The FlowPot system is based on autoclaved and washed soil, and thus contains organic carbon; however, in this system, the nutrient content cannot be controlled [7^{**},29^{**}]. The choice of systems should therefore be based on the ultimate aim of the experiment.

Reconstituting root microbiomes using SynComs

Using cultivated pure strains, scientists are able to constitute microbial SynComs in axenic systems based on the natural microbiome patterns, obtained from the correlation between microbes and plant phenotype [10^{**}], the comparison of root microbiomes between wild-type and mutant plants [43^{**}], or network analyses [7^{**}] (Figure 2e). These

approaches provide new opportunities to validate the root microbiome assembly patterns observed in natural soil conditions, study the effect of the root microbiome on plant fitness under different environmental conditions, and investigate microbe–microbe interactions and microbial gene functions [44,45,46] (Figure 2f).

Using SynComs in axenic conditions provides researchers with a controlled ecological environment in which microbes, nutrients, and plant genotypes are strictly monitored, generating reproducible results that can be used to validate observations of root microbiome patterns in natural soils. For example, Lebeis *et al.* detected alterations in the *A. thaliana* root microbiomes of salicylic acid signaling mutants compared with the wild-type plants grown in natural soils or in a calcined clay-based reconstitution system containing 38 bacterial strains [43**]. Four of the seven bacterial strains showing a difference in abundance between wild-type and mutant plants in the SynCom experiments belonged to families that showed differential abundance between wild-type and mutants in natural soils. Based on the evidence obtained from natural soil and synthetic laboratory conditions, the authors therefore concluded that the salicylic acid-mediated plant innate immunity signaling pathways modulate the assembly of the *Arabidopsis* root microbiome at the family level [43**]. A similar logic was applied by Castrillo *et al.*, who investigated the effect of the phosphate starvation pathways on the establishment of the *Arabidopsis* root microbiome [40**,41]. These authors identified altered root microbiome compositions between wild-type plants and several phosphate starvation mutants in both natural soils and SynCom experiments using Murashige and Skoog agar plates.

In addition, SynCom approaches using sequenced cultivated bacteria provide simple and reproducible systems to study the role of root metabolites (e.g. triterpenes, aromatic acids, coumarin, and camalexin) on the interactions between the root microbiome and the host plants in complex natural soils [47**,48,49*,50,51**]. In summary, experimentation using SynComs under controlled conditions permits the validation of the assembly patterns of the root microbiome and complements studies of the root microbiome in natural soil conditions.

A major benefit of inoculating germ-free plants with SynCom in axenic systems is the ability to study the function of the root microbiome under different environmental conditions. For example, Duran *et al.* inoculated germ-free *Arabidopsis* plants in the soil-based FlowPot system with different SynComs, including bacteria, fungi, and oomycetes, and found that the root bacterial microbiome protects *Arabidopsis* plants against filamentous eukaryotes (fungi and oomycetes) in natural soils [7**]. Castrillo *et al.* inoculated germ-free plants with SynComs under low-phosphate conditions, revealing that bacterial

inoculation dramatically increased the plant phosphate starvation responses, and that master regulators of phosphate starvation pathways directly suppress of the plant innate immune system [40**]. In addition, the characterization of a small set of SynComs proved to be sufficient for predicting the effect of the microbiome on the phosphate utilization by *Arabidopsis* [41]. A recent follow-up study revealed that *Burkholderia* competed with plants for phosphate utilization, exacerbating plant phosphate starvation [42]. Using bacteria cultivated from the microbiomes of *indica* and *japonica* rice varieties, Zhang *et al.* revealed that the rice root microbiome facilitates the utilization of organic nitrogen by the plants [8**]. Based on correlation analysis of root microbiomes in wild disease-resistant and susceptible tomato cultivars, Kwak *et al.* successfully cultivated a target flavobacterium and demonstrated that it promoted wilt resistance when inoculated into tomato roots [10**]. Genome analyses of cultivated *Pseudomonas* led to the identification of lipopeptide biosynthesis and quorum-sensing as the major genomic differences between pathogenic and commensal root bacteria [46], in addition to the genes required for root bacteria to evade plant defense or induce plant susceptibility [44,45]. These examples demonstrate the utility of SynComs or cultivated microbes in elucidating the effect of the root microbiome on plant fitness under different environmental conditions.

Cultivated microbes and SynCom approaches provide unique opportunities for the exploration of microbe–microbe interactions and microbial gene functions. Using an agar system and a SynCom consisting of seven bacterial strains cultivated from maize roots, Niu *et al.* showed that the removal of *Enterobacter cloacae* resulted in the complete loss of microbial community structure in the roots of maize seedlings, demonstrating the importance of microbe–microbe interactions and the existence of keystone species in the microbiome [52]. Duran and colleagues performed a large-scale binary assay between cultivated root-derived bacteria, fungi, and oomycetes, revealing that the biocontrol activity of bacteria against filamentous eukaryotes is a redundant trait that maintains the interkingdom balance of the root microbiome [7**]. Levy *et al.* found that the survival of *Escherichia coli* and six leaf bacterial isolates was reduced 10^2 – 10^6 fold after being incubated with wild-type *Acidovorax* compared with the delta-*HydE1* and Type VI secretion system (T6SS) *Acidovorax* mutants, suggesting that Hyde and T6SS-mediated antimicrobial activity could be used to manipulate the microbiome [21**]. These studies reveal the value of cultivated microbes in dissecting microbe–microbe interactions.

Conclusions

Because of the complexity of soil environments, current SynCom approaches can only partially mimic natural conditions. The low complexity of SynCom systems may mean that some important microbiome members

or nutrients are lacking; however, microbial cultivation followed by experimentation using SynComs in axenic systems permits us to elucidate the causality between the root microbiome and plant phenotypes, and to dissect the interactions between microbiome members under natural soil conditions. More cultivated microbes, genomes, improved axenic systems, and the systematic investigation of different functional microbiome members are needed to further reveal the functions and mechanisms of the root microbiome in plant growth and health.

Acknowledgements

This work was supported by the Strategic Priority Research Program of the Chinese Academy of Sciences (grant nos. XDB11020700), the Key Research Program of Frontier Sciences of the Chinese Academy of Science (grant nos. QYZDB-SSW-SMC021), the National Natural Science Foundation of China (grant nos. 31772400), and the Key Research Program of the Chinese Academy of Sciences (grant nos. KFZD-SW-219).

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
 - of outstanding interest
1. Bulgarelli D, Rott M, Schlaeppi K, Ver Loren van Themaat E, •• Ahmadinejad N, Assenza F, Rauf P, Huettel B, Reinhardt R, Schmelzer E *et al.*: **Revealing structure and assembly cues for *Arabidopsis* root-inhabiting bacterial microbiota.** *Nature* 2012, **488**:91-95
Defined the distinctive structure of bacterial microbiota in *Arabidopsis* roots comparing with unplanted natural soils revealed that root microbiota are determined by soil types and plant genotypes.
 2. Jansson JK, Hofmockel KS: **The soil microbiome—from metagenomics to metaphenomics.** *Curr Opin Microbiol* 2018, **43**:162-168.
 3. Lundberg DS, Lebeis SL, Paredes SH, Yourstone S, Gehring J, •• Malfatti S, Tremblay J, Engelbrektson A, Kunin V, d Rio TG *et al.*: **Defining the core *Arabidopsis thaliana* root microbiome.** *Nature* 2012, **488**:86
Defined the distinctive structure of bacterial microbiota in *Arabidopsis* roots comparing with unplanted natural soils; revealed that root microbiota are determined by soil types and plant genotypes.
 4. Zhang J, Zhang N, Liu Y-X, Zhang X, Hu B, Qin Y, Xu H, Wang H, Guo X, Qian J *et al.*: **Root microbiota shift in rice correlates with resident time in the field and developmental stage.** *Sci China Life Sci* 2018, **61**:613-621.
 5. Edwards JA, Santos-Medellin CM, Liechty ZS, Nguyen B, Lurie E, Eason S, Phillips G, Sundaresan V: **Compositional shifts in root-associated bacterial and archaeal microbiota track the plant life cycle in field-grown rice.** *PLoS Biol* 2018, **16**:e2003862.
 6. Lu T, Ke M, Lavoie M, Jin Y, Fan X, Zhang Z, Fu Z, Sun L, Gillings M, Peñuelas J *et al.*: **Rhizosphere microorganisms can influence the timing of plant flowering.** *Microbiome* 2018, **6**:231.
 7. Durán P, Thiergart T, Garrido-Oter R, Agler M, Kemen E, Schulze-Lefert P, Hacquard S: **Microbial interkingdom interactions in roots promote *Arabidopsis* survival.** *Cell* 2018, **175**:973-983. e914
Revealed the function of root bacteria to protect *Arabidopsis* growth against soil-borne fungi and oomycetes in natural soil.
 8. Zhang J, Liu Y-X, Zhang N, Hu B, Jin T, Xu H, Qin Y, Yan P, •• Zhang X, Guo X *et al.*: ***NRT1.1B* is associated with root microbiota composition and nitrogen use in field-grown rice.** *Nat Biotechnol* 2019, **37**:676-684
Revealed that rice plants recruit root microbiota to utilize soil nitrogen.
 9. Hiruma K, Gerlach N, Sacristán S, Nakano Ryohei T, Hacquard S, •• Kracher B, Neumann U, Ramírez D, Bucher M, O'Connell Richard J *et al.*: **Root endophyte colleotrichum tofieldiae confers plant fitness benefits that are phosphate status dependent.** *Cell* 2016, **165**:464-474
Identified the symbiotic interaction between endophytic fungus *C. tofieldiae* and *Arabidopsis* for Pi utilization.
 10. Kwak M-J, Kong HG, Choi K, Kwon S-K, Song JY, Lee J, Lee PA, •• Choi SY, Seo M, Lee HJ *et al.*: **Rhizosphere microbiome structure alters to enable wilt resistance in tomato.** *Nat Biotechnol* 2018, **36**:1100
Revealed that rhizosphere microbiota contribute to wilt resistance in tomato. Metagenome was used for targeted bacteria isolation.
 11. Berendsen RL, Vismans G, Yu K, Song Y, de Jonge R, Burgman WP, Burmölle M, Herschend J, Bakker PAHM, Pieterse CMJ: **Disease-induced assemblage of a plant-beneficial bacterial consortium.** *ISME J* 2018, **12**:1496-1507.
 12. Hu J, Wei Z, Friman V-P, Gu S-H, Wang X-F, Eisenhauer N, Yang T-J, Ma J, Shen Q-R, Xu Y-C *et al.*: **Probiotic diversity enhances rhizosphere microbiome function and plant disease suppression.** *mBio* 2016, **7**:e01790-01716.
 13. Hu L, Robert CAM, Cadot S, Zhang X, Ye M, Li B, Manzo D, • Chervet N, Steinger T, van der Heijden MGA *et al.*: **Root exudate metabolites drive plant-soil feedbacks on growth and defense by shaping the rhizosphere microbiota.** *Nat Commun* 2018, **9**:2738
Illustrated the function root exudates to shape root microbiota for defense in maize.
 14. Beckers B, Op De Beeck M, Weyens N, Boerjan W, Vangronsveld J: **Structural variability and niche differentiation in the rhizosphere and endosphere bacterial microbiome of field-grown poplar trees.** *Microbiome* 2017, **5**:25.
 15. Edwards J, Johnson C, Santos-Medellin C, Lurie E, Podishetty NK, •• Bhatnagar S, Eisen JA, Sundaresan V: **Structure, variation, and assembly of the root-associated microbiomes of rice.** *Proc Natl Acad Sci U S A* 2015, **112**:E911-E920
Provided evidence for rice root microbiota assembly from soil to root by time-course data in the greenhouse.
 16. Shi Y, Li Y, Xiang X, Sun R, Yang T, He D, Zhang K, Ni Y, Zhu Y-G, Adams JM *et al.*: **Spatial scale affects the relative role of stochasticity versus determinism in soil bacterial communities in wheat fields across the North China Plain.** *Microbiome* 2018, **6**:27.
 17. Lee SA, Park J, Chu B, Kim JM, Joa J-H, Sang MK, Song J, Weon H-Y: **Comparative analysis of bacterial diversity in the rhizosphere of tomato by culture-dependent and -independent approaches.** *J Microbiol* 2016, **54**:823-831.
 18. Hartman K, van der Heijden MGA, Roussely-Provent V, Walser J-C, Schlaeppi K: **Deciphering composition and function of the root microbiome of a legume plant.** *Microbiome* 2017, **5**:2.
 19. Walters WA, Jin Z, Youngblut N, Wallace JG, Sutter J, Zhang W, González-Peña A, Peiffer J, Koren O, Shi Q *et al.*: **Large-scale replicated field study of maize rhizosphere identifies heritable microbes.** *Proc Natl Acad Sci U S A* 2018, **115**:7368.
 20. Bai Y, Müller DB, Srinivas G, Garrido-Oter R, Potthoff E, Rott M, •• Dombrowski N, Münch PC, Spaepen S, Remus-Emsermann M *et al.*: **Functional overlap of the *Arabidopsis* leaf and root microbiota.** *Nature* 2015, **528**:364-369
Established systematic bacterial culture collection in *Arabidopsis* roots and leaves, which paves the way for functional and mechanistic studies of plant microbiota.
 21. Levy A, Salas Gonzalez I, Mittelviehhaus M, Clingenpeel S, Herrera •• Paredes S, Miao J, Wang K, Devescovi G, Stillman K, Monteiro F *et al.*: **Genomic features of bacterial adaptation to plants.** *Nat Genet* 2018, **50**:138-150
Identified the key bacterial genes for colonization in roots using the genomes of cultivated bacteria.
 22. Armanhi JSL, de Souza RSC, Damasceno Nd B, de Araújo LM, Imperial J, Arruda P: **A community-based culture collection for**

- targeting novel plant growth-promoting bacteria from the sugarcane microbiome.** *Front Plant Sci* 2018;8.
23. Khan Chowdhury MDE, Jeon J, Ok Rim S, Park Y-H, Kyu Lee S, Bae H: **Composition, diversity and bioactivity of culturable bacterial endophytes in mountain-cultivated ginseng in Korea.** *Sci Rep* 2017, **7**:10098.
24. Vorholt JA, Vogel C, Carlström CI, Müller DB: **Establishing causality: opportunities of synthetic communities for plant microbiome research.** *Cell Host Microbe* 2017, **22**:142-155
Discussed to establish causality using SynCom.
25. Schlaeppi K, Dombrowski N, Oter RG, Ver Loren van Themaat E, Schulze-Lefert P: **Quantitative divergence of the bacterial root microbiota in *Arabidopsis thaliana* relatives.** *Proc Natl Acad Sci U S A* 2014, **111**:585-592
Reveal the divergence and conservation of root microbiota in *Arabidopsis* and relatives.
26. Agler MT, Ruhe J, Kroll S, Morhenn C, Kim S-T, Weigel D, Kemen EM: **Microbial hub taxa link host and abiotic factors to plant microbiome variation.** *PLoS Biol* 2016, **14**:e1002352
Identified hub bacterial taxa in the complex interactions among bacteria, fungi, oomycetes and *Arabidopsis* leaves.
27. Lebeis SL, Rott M, Dangl JL, Schulze-Lefert P: **Culturing a plant microbiome community at the cross-Rhodes.** *New Phytol* 2012, **196**:341-344.
28. Zamioudis C, Korteland J, Van Pelt JA, van Hamersveld M, Dombrowski N, Bai Y, Hanson J, Van Verk MC, Ling H-Q, Schulze-Lefert P *et al.*: **Rhizobacterial volatiles and photosynthesis-related signals coordinate MYB72 expression in *Arabidopsis* roots during onset of induced systemic resistance and iron-deficiency responses.** *Plant J* 2015, **84**:309-322
Revealed root bacteria's function in Fe uptake.
29. Kremer JM, Paasch BC, Rhodes D, Thireault C, Froehlich JE, Schulze-Lefert P, Tiedje JM, He SY: **FlowPot axenic plant growth system for microbiota research.** *bioRxiv* 2018:254953
Established the soil-based germ free system.
30. Zou Y, Xue W, Luo G, Deng Z, Qin P, Guo R, Sun H, Xia Y, Liang S, Dai Y *et al.*: **1520 reference genomes from cultivated human gut bacteria enable functional microbiome analyses.** *Nat Biotechnol* 2019, **37**:179-520185.
31. Oberhardt MA, Zarecki R, Gronow S, Lang E, Klenk H-P, Gophna U, Ruppin E: **Harnessing the landscape of microbial culture media to predict new organism-media pairings.** *Nat Commun* 2015, **6**:8493.
32. Ward T, Larson J, Meulemans J, Hillmann B, Lynch J, Sidiropoulos D, Spear JR, Caporaso G, Blekman R, Knight R *et al.*: **BugBase predicts organism-level microbiome phenotypes.** *bioRxiv* 2017:133462.
33. Lombard V, Golaconda Ramulu H, Drula E, Coutinho PM, Henrissat B: **The carbohydrate-active enzymes database (CAZy) in 2013.** *Nucleic Acids Res* 2013, **42**:D490-D495.
34. Jia B, Raphenya AR, Alcock B, Waglechner N, Guo P, Tsang KK, Lago BA, Dave BM, Pereira S, Sharma AN *et al.*: **CARD2017: expansion and model-centric curation of the comprehensive antibiotic resistance database.** *Nucleic Acids Res* 2017, **45**:D566-D573.
35. Kato S, Yamagishi A, Daimon S, Kawasaki K, Tamaki H, Kitagawa W, Abe A, Tanaka M, Sone T, Asano K *et al.*: **Isolation of previously uncultured slow-growing bacteria by using a simple modification in the preparation of agar media.** *Appl Environ Microbiol* 2018, **84**:e00807-00818.
36. Edgar RC, Flyvbjerg H: **Error filtering, pair assembly and error correction for next-generation sequencing reads.** *Bioinformatics* 2015, **31**:3476-3482.
37. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP: **DADA2: High-resolution sample inference from Illumina amplicon data.** *Nat Methods* 2016, **13**:581.
38. Amir A, McDonald DP, Navas-Molina JA, Kopylova E, Morton JT, Zech Xu Z, Kightley EP, Thompson LR, Hyde ER, Gonzalez A *et al.*: **Deblur rapidly resolves single-nucleotide community sequence patterns.** *mSystems* 2017, **2**:e00191-00116.
39. Karasov TL, Almaro J, Friedemann C, Ding W, Giolai M, Heavens D, Kersten S, Lundberg DS, Neumann M, Regalado J *et al.*: ***Arabidopsis thaliana* and *Pseudomonas* pathogens exhibit stable associations over evolutionary timescales.** *Cell Host Microbe* 2018, **24**:168-179.e164
Revealed evolutionary traits of *Pseudomonas* associated with *Arabidopsis* leaves using the genomes of large scale cultivated strains.
40. Castrillo G, Teixeira PJPL, Paredes SH, Law TF, de Lorenzo L, Feltcher ME, Finkel OM, Breakfield NW, Mieczkowski P, Jones CD *et al.*: **Root microbiota drive direct integration of phosphate stress and immunity.** *Nature* 2017, **543**:513
Revealed that root microbiota integrate phosphate stress signaling and plant immunity using 16S rRNA gene sequencing and SynComs.
41. Herrera Paredes S, Gao T, Law TF, Finkel OM, Mucyn T, Teixeira PJPL, Salas González I, Feltcher ME, Powers MJ, Shank EA *et al.*: **Design of synthetic bacterial communities for predictable plant phenotypes.** *PLoS Biol* 2018, **16**:e2003962.
42. Finkel OM, Salas-González I, Castrillo G, Spaepen S, Law TF, Jones CD, Dangl JL: **The effects of soil phosphorous content on microbiota are driven by the plant phosphate starvation response.** *bioRxiv* 2019:608133.
43. Lebeis SL, Paredes SH, Lundberg DS, Breakfield N, Gehring J, McDonald M, Malfatti S, Glavina del Rio T, Jones CD, Tringe SG *et al.*: **Salicylic acid modulates colonization of the root microbiome by specific bacterial taxa.** *Science* 2015, **349**:860-864
The first paper to use the term 'SynCom'. Revealed the function of SA-mediation innate immune system to modulate *Arabidopsis* root microbiota by 16S rRNA gene sequencing and SynComs.
44. Liu Z, Beskrvnaya P, Melnyk RA, Hossain SS, Khorasani S, O'Sullivan LR, Wiesmann CL, Bush J, Richard JD, Haney CH: **A genome-wide screen identifies genes in rhizosphere-associated *Pseudomonas* required to evade plant defenses.** *mBio* 2018, **9**:e00433-00418.
45. Melnyk RA, Beskrvnaya P, Liu Z, Song Y, Haney CH: **Bacterially produced spermidine induces plant systemic susceptibility to pathogens.** *bioRxiv* 2019:517870.
46. Melnyk RA, Hossain SS, Haney CH: **Convergent gain and loss of genomic islands drive lifestyle changes in plant-associated *Pseudomonas*.** *ISME J* 2019, **13**:1575-1588.
47. Huang AC, Jiang T, Liu Y-X, Bai Y-C, Reed J, Qu B, Goossens A, Nützmann H-W, Bai Y, Osbourn A: **A specialized metabolic network selectively modulates *Arabidopsis* root microbiota.** *Science* 2019, **364**:eaau6389
Revealed *Arabidopsis*-specialized triterpenes selective modulate root microbiota using 16S rRNA gene sequencing and cultivated strains.
48. Stringlis IA, Yu K, Feussner K, de Jonge R, Van Bentum S, Van Verk MC, Berendsen RL, Bakker PAHM, Feussner I, Pieterse CMJ: **MYB72-dependent coumarin exudation shapes root microbiome assembly to promote plant health.** *Proc Natl Acad Sci U S A* 2018, **115**:E5213-E5222.
49. Voges MJEEE, Bai Y, Schulze-Lefert P, Sattely ES: **Plant-derived coumarins shape the composition of an *Arabidopsis* synthetic root microbiome.** *Proc Natl Acad Sci U S A* 2019:201820691
Revealed that plant-derived coumarins regulate root pseudomonas strain in low Fe condition using SynComs.
50. Koprivova A, Schuck S, Jacoby RP, Klinkhammer I, Welter B, Leson L, Martyn A, Nauen J, Grabenhorst N, Mandelkow JF *et al.*: **Root-specific camalexin biosynthesis controls the plant growth-promoting effects of multiple bacterial strains.** *Proc Natl Acad Sci U S A* 2019, **116**:15735-15744.
51. Zhalnina K, Louie KB, Hao Z, Mansoori N, da Rocha UN, Shi S, Cho H, Karaoz U, Loque D, Bowen BP *et al.*: **Dynamic root exudate chemistry and microbial substrate preferences drive patterns in rhizosphere microbial community assembly.** *Nat Microbiol* 2018, **3**:470-480
Revealed the function of microbial preferences for roots exudates in root microbiota assembly.
52. Niu B, Paulson JN, Zheng X, Kolter R: **Simplified and representative bacterial community of maize roots.** *Proc Natl Acad Sci U S A* 2017, **114**:E2450.